

Flagyl® Cures Trichomonal Infection in Both

brand of metronidazole

Although *Trichomonas vaginalis* infection occurs in only 5 to 10 per cent* of men, careful diagnosis will demonstrate the condition in about half of all husbands of infected women. Nine investigators* reported an average incidence of 50.8 per cent in exposed consorts.

Many clinicians have achieved a high degree of success in treating trichomonal vaginitis only after they have recognized the importance of sexual partners in perpetuating the infection. Crowley* has asserted, "it was not until we acted on this key premise that we were able to obtain positive and lasting results in our management of recurrent vaginal trichomoniasis."

Simple ten-day oral treatment with Flagyl virtually assures elimination of established trichomonal infection in men. In twenty-two of twenty-seven studies* data on the results of treating male patients revealed that all men treated with Flagyl were cured.

Indications: Flagyl is indicated in the treatment of trichomoniasis in both men and women.

Contraindications: Pregnancy; disease of the central nervous system; evidence or history of blood dyscrasia.

Precaution: Complete blood cell counts should be made before, during and after therapy, especially if a second course is necessary.

Side effects: Infrequent and minor side effects include nausea, metallic taste and furry tongue. Gastrointestinal disturbances, flushing and headache sometimes occur, especially with concomitant ingestion of alcohol. The taste of alcoholic beverages may be altered. Other effects, all reported in an incidence of less than 1 per cent, are diarrhea, dizziness, vaginal dryness and burning, dry mouth, rash, urticaria, gastritis, drowsiness, insomnia, pruritus, sore tongue, darkened urine, anorexia, vomiting, epigastric distress, dysuria, depression, vertigo, incoordination, ataxia, abdominal cramping, constipation, stomatitis, numbness or paresthesia of an extremity, joint pains, confusion, irritability, weakness, cystitis, pelvic pressure, dyspareunia, fever, polyuria, incontinence, decreased libido, nasal congestion, proctitis and pyuria. Elimination of trichomonads may aggravate candidiasis.

Dosage and Administration: *In women:* one 250-mg. oral tablet three times daily for ten days. A vaginal insert of 500 mg. is available for local therapy when desired. When used, one vaginal insert should be placed high in the vaginal vault each day for ten days; concurrently two oral tablets should be taken daily.

In men: When trichomonads are demonstrated, one 250-mg. oral tablet twice daily for ten days in conjunction with treatment of his female partner.

Dosage Forms: Oral tablets—250 mg.

Vaginal inserts—500 mg.

*Complete list of references on request.

Your assistant needs all the help she can get.

And she'll get it. From Blue Shield.

A medical assistant's work is never done. There are endless bills to pay, forms to fill out, messages to take, forms to fill out, patients to see, and more forms to fill out.

Often it's hard to keep facts straight and things in order. Even the best assistants need a little help once in a while. That's why Blue Shield has set up an assistant's program designed by key medical assistants and medical assistants' groups who know the problems.

For instance, if your assistant gets stuck with a problem concerning a patient's coverage or claims, paying bills or keeping books, there's a Blue Shield representative just a phone call away (212) 340-5131. And through our frequent training courses, special publications and her own medical assistant's manual, she'll be able to save you time and paperwork.

The more your assistant knows about her job, the easier yours will be. And Blue Shield will make sure she gets all the help she'll need.



GREATER NEW YORK'S

BLUE SHIELD

UNITED MEDICAL SERVICE, INC.

The Bulletin of The New York Academy of Medicine, Vol. 45, No. 6, June, 1969. Published monthly by The New York Academy of Medicine, 2 East 103 Street, New York, N. Y. 10029. Entered as second-class matter February 3, 1928, at the Post Office at New York, N. Y., under the act of August 24, 1912. Postage paid at New York, N. Y. Annual subscription United States \$10.00. Canada \$11.00. All other countries \$12.00. Single copies \$2.00.



Scaled for the patient with high-level anxiety

Librium® (chlordiazepoxide HCl) 25-mg capsules

Because anxiety varies widely from patient to patient, and even in the same individual, Librium (chlordiazepoxide HCl) is supplied in various dosage strengths to suit the level of anxiety. Thus, during periods of acute emotional stress, the patient may need 25 mg Librium *t.i.d.* for relief. In mild to moderate anxiety, smaller doses of 5 or 10 mg, given three or four times daily, usually suffice.

The resulting improvement in outlook is a characteristic benefit of Librium therapy, utilized as an adjunct to your counsel and reassurance. Another advantage: Librium may also be used concomitantly with certain specific medications of other classes of drugs, whenever anxiety is a significant component of the clinical profile.

Before prescribing, please consult complete product information, a summary of which follows:

Indications: Indicated when anxiety, tension and apprehension are significant components of the clinical profile.

Contraindications: Patients with known hypersensitivity to the drug.

Warnings: Caution patients about possible combined effects with alcohol and other CNS depressants. As with all CNS-acting drugs, caution patients against hazardous occupations requiring com-

plete mental alertness (e.g., operating machinery, driving). Though physical and psychological dependence have rarely been reported on recommended doses, use caution in administering to addiction-prone individuals or those who might increase dosage; withdrawal symptoms (including convulsions), following discontinuation of the drug and similar to those seen with barbiturates, have been reported. Use of any drug in pregnancy, lactation, or in women of childbearing age requires that its potential benefits be weighed against its possible hazards.

Precautions: In the elderly and debilitated, and in children over six, limit to smallest effective dosage (initially 10 mg or less per day) to preclude ataxia or oversedation, increasing gradually as needed and tolerated. Not recommended in children under six. Though generally not recommended, if combination therapy with other psychotropics seems indicated, carefully consider individual pharmacologic effects, particularly in use of potentiating drugs such as MAO inhibitors and phenothiazines. Observe usual precautions in presence of impaired renal or hepatic function. Paradoxical reactions (e.g., excitement, stimulation and acute rage) have been reported in psychiatric patients and hyperactive aggressive children. Employ usual precautions in treatment of anxiety states with evidence of impending depression; suicidal tendencies may be present and protective measures necessary. Variable effects on blood coagulation have been reported very rarely in patients receiving the drug and oral anticoagulants; causal relationship has not been established clinically.

Adverse Reactions: Drowsiness, ataxia and confusion may occur, especially in the elderly and debilitated. These are

reversible in most instances by proper dosage adjustment, but are also occasionally observed at the lower dosage ranges. In a few instances syncope has been reported. Also encountered are isolated instances of skin eruptions, edema, minor menstrual irregularities, nausea and constipation, extrapyramidal symptoms, increased and decreased libido—all infrequent and generally controlled with dosage reduction; changes in EEG patterns (low-voltage fast activity) may appear during and after treatment; blood dyscrasias (including agranulocytosis), jaundice and hepatic dysfunction have been reported occasionally, making periodic blood counts and liver function tests advisable during protracted therapy.

Usual Daily Dosage: Individualize for maximum beneficial effects. *Oral*—

Adults: Mild and moderate anxiety and tension, 5 or 10 mg *t.i.d.* or *q.i.d.*; severe states, 20 or 25 mg *t.i.d.* or *q.i.d.* Geriatric patients: 5 mg *b.i.d.* to *q.i.d.* (See **Precautions**.)

Supplied: Librium®(chlordiazepoxide HCl) Capsules, 5 mg, 10 mg and 25 mg—bottles of 50. Libritabs®(chlordiazepoxide) Tablets, 5 mg, 10 mg and 25 mg—bottles of 100. With respect to clinical activity, capsules and tablets are indistinguishable.

when tablets are preferred:


Libritabs®
(chlordiazepoxide)



Roche
LABORATORIES

Division of Hoffmann-La Roche Inc.
Nutley, New Jersey 07110

now for your non-hospitalized patients



Serum Chemistry Graph provided is 8" x 11"

blood chemistry profiles

run on
TECHNICON®
Sequential Multiple Analyzer

only \$15⁰⁰

FOR A ~~12~~-TEST BLOOD CHEMISTRY PROFILE

Heretofore available only to patients in a few large hospitals, PATHOLOGICAL LABORATORY now provides this rapid automated analytical service for the non-hospitalized patient.

A Serum Chemistry Graph of each analysis summarizes in graphic form the results of all 12 tests together, superimposed over a background of normal values.

Patients may be directed to the laboratory; or blood samples can be mailed in post-paid containers provided by our laboratory. The Serum Chemistry Graph and a report in the usual numerical format will be mailed the same day samples are received. Free messenger pick-up is available in select areas.

PATHOLOGICAL LABORATORY

40-03 NATIONAL ST., CORONA, N.Y. 11368

TELEPHONE 212-639-5403

**send for
post-paid
blood sample
mailing
containers**

PATHOLOGICAL LABORATORY • 40-03 National St., Corona, N.Y. 11368

Gentlemen:

- ☐ Please send additional literature
- ☐ Please send post-paid containers for blood samples

Dr.

Street

City State Zip

*“For all the happiness
mankind can gain
It is not in pleasure,
but in rest from pain.”*

John Dryden

‘Empirin’ Compound with Codeine Phosphate gr. 1/2 No. 3


Each tablet contains: Codeine Phosphate gr. 1/2 (Warning—May be habit forming), Phenacetin gr. 2 1/2, Aspirin gr. 3 1/2, Caffeine gr. 1/2.

gives your patient rest from pain

‘B.W. & Co.’ narcotic products are Class “B”, and as such are available on oral prescription, where State law permits.



BURROUGHS WELLCOME & CO. (U.S.A.) INC., Tuckahoe, N. Y.



Darvon[®] Compound-65

Each Pulvule[®] contains 65 mg. propoxyphene hydrochloride, 227 mg. aspirin, 162 mg. phenacetin, and 32.4 mg. caffeine.

Lilly

Additional information
available upon request.
Eli Lilly and Company
Indianapolis, Indiana 46206

Over 5 million prescriptions already filled at Macy's

*All 13 Macy Prescription Centres
satisfy your patients with
efficient professional service*



- You may phone in your patients' prescriptions (except for narcotics, amphetamines, and barbiturates).
- Your patients may charge their prescriptions on their Macy Shopping Accounts or C-T Plans.
- Your patients may have their prescriptions delivered anywhere free of extra charge.
- We serve your patient with a full line of convalescent and diabetic needs.

Macy's Herald Square, OX 5-4400
Macy's Parkchester, TA 8-7000
Macy's Jamaica, OL 7-9000
Macy's Flatbush, UL 6-5000
Macy's Roosevelt Field, PI 6-8200
Macy's Huntington, AR 1-3000
Macy's White Plains, WH 6-5015
Macy's New Haven, Conn. 624-9271
Macy's Bay Shore, MO 5-8400
Macy's Queens, AR 1-9100
Macy's Colonie, 459-1950
Macy's New Rochelle, 633-7700
Macy's Smith Haven, 724-4900

BULLETIN of THE NEW YORK ACADEMY OF MEDICINE

The *Bulletin* contains scientific papers presented at the Academy, including:

- ADDRESSES PRESENTED AT STATED AND ANNUAL MEETINGS
- PAPERS PRESENTED AT SECTIONS OF THE ACADEMY
- PROCEEDINGS OF SYMPOSIA AND CONFERENCES
- TELEVISED CLINICAL SCIENCE SEMINARS
- PUBLIC HEALTH REPORTS
- NEW YORK PATHOLOGICAL SOCIETY—ABSTRACTS

Published monthly

Subscription office: S-H Service Agency, Inc.
31 East 10th Street, New York, N. Y. 10003

**Annual subscription: United States \$10.00, Canada \$11.00,
all other countries \$12.00 — single copies \$2.00**

Three important titles from **HAFNER PUBLISHING COMPANY**

DOWNEY, HAROLD**Handbook of Hematology**

4 vols. 3136 pages, colored plates, illus.
(1938) Reprint 1965 \$137.50

"These four volumes contain a critical digest of the international literature of hematology as well as conclusions and opinions of the authors, based on personal research in their various fields. The illustrations, particularly those done in color, are an important feature, as are the bibliographies appended to most of the sections. This work is intended for clinicians, pathologists, histologists, immunologists, pediatricians etc." (Hawkins)

"A publishing event of 1965 was the reissue of Downey's classic four volume work." (JAMA, April 4, 1966)

COWDRY, E. V.

Special Cytology—The form and functions of the cell in health and disease.

2nd ed. 3 vols. 757 illus. (1932)
Reprint 1963 \$52.50

This work has become a classic, covering literature up to 1932 as no other book in experimental morphology does. It serves as a base line from which most modern work has evolved.

DANIELLI, J. F.**Cell Physiology and Pharmacology**

viii, 156 pages, 3 plates, 21 illus., 22 tables. (1950) Reprint 1968 \$8.00

The author, an honorary lecturer in Pharmacology at University College, London, specialized in problems which are encountered when searching for a new drug. This book was intended to direct attention to the biological aspects of drug action and to indicate some of the more important factors, on the cellular level, which must be considered by students of drug action. This very valuable contribution should be of great value for everyone concerned with these problems—pharmacologists, chemists, biologists, medical men, and the like.

Send your orders to **STECHERT-HAFNER, INC.**
31 East 10th Street, New York, N. Y. 10003

ADVERTISERS INDEX

Advertisers	Page
Bull. NYAM	VIII
Burroughs Wellcome & Co., Inc.	VI
Group Health Insurance	3rd cover
Eli Lilly & Co.	VII
Macy Prescription Centres	VIII
Pathological Laboratory	V
Physicians Wanted	IX
Roche Laboratories	III, IV, 4th cover
Searle Co., G. D.	2nd cover
Stechert-Hafner, Inc.	IX
United Medical Service, Inc.	II

PHYSICIANS WANTED


The State of Virginia has opportunities for Physicians in Public Health. At the present time, nine vacancies exist in varied locations for physicians, who as State employees would be assigned to direct public health and medical care programs in local health departments. Programs include traditional public health services, plus activities under Medicare and Medicaid. In-service training and opportunity for post graduate study provided. Liberal fringe benefits.

Qualifications: American citizen or papers, eligibility to practice medicine in Virginia, two years experience in civilian or military practice, good speaking ability with the English language, age under 55 (under 60 if MPH). Salary \$17,900, with regular increments to \$21,400 in five years. \$19,600 to \$24,500 in two districts which require MPH. Note that living expenses are less in the South, and we are not crowded.

For further information, inquire Director, Local Health Services, Virginia State Department of Health, Richmond, Virginia 23219. Better still, stop by and see us when traveling South. We are three blocks off I-95 (Broad Street Exit, West), in downtown Richmond, James Madison Office Building, 5th Floor.

GREAT AMERICANS WHO WERE DOCTORS...

... NUMBER SIX OF A SERIES



John Kearsley Mitchell
(1792-1858)

After sailing on three voyages to China and the East Indies as ship's surgeon, Dr. Mitchell set up practice in Philadelphia and joined the faculty of the Philadelphia Medical Institute. He is known primarily as the author of "Saint Helena, a Poem by a Yankee," and "Indecision, a Tale of the Far West and Other Poems."

We AT GHI know little about the writing of poetry. But when it comes to writing contracts that protect over one million New Yorkers, we are professionals. We ask you to join us as a Participating Family Doctor or a Participating Physician. This will be a reciprocally satisfactory relationship. For you, there will be the feeling of accomplishment that comes with belonging to a group that extends quality medical care to many thousands of people. For us, there will be a strengthening of the "service-benefit" feature that is at the core of our existence.

Join the approximately 10,000 doctors who now participate in GHI's Plans. Write to us today for more information about why and how you should participate in GHI. Do it now — while you're thinking about it!



Group Health Insurance, Inc. / 227 West 40th Street, New York, N.Y. 10018 / Phone: 564-8900

Animal Pharmacology: Calming and taming the monkey

Before



After



Naturally vicious laboratory monkeys show consistently

aggressive behavior toward their handlers. On Valium®(diazepam),

they become quite "tame," yet remain alert and coordinated.

Before prescribing, please consult complete product information, a summary of which follows:

Indications: Tension and anxiety states; somatic complaints which are concomitants of emotional factors; psychoneurotic states manifested by tension, anxiety, apprehension, fatigue, depressive symptoms or agitation; acute agitation, tremor, delirium tremens and hallucinosis due to acute alcohol withdrawal; adjunctively in skeletal muscle spasm due to reflex spasm to local pathology, spasticity caused by upper motor neuron disorders, athetosis, stiff-man syndrome, convulsive disorders (not for sole therapy).

Contraindicated: Known hypersensitivity to the drug. Children under 6 months of age. Acute narrow angle glaucoma.

Warnings: Not of value in psychotic patients. Caution against hazardous occupations requiring complete mental alertness. When used adjunctively in convulsive dis-

orders, possibility of increase in frequency and/or severity of grand mal seizures may require increased dosage of standard anticonvulsant medication; abrupt withdrawal may be associated with temporary increase in frequency and/or severity of seizures. Advise against simultaneous ingestion of alcohol and other CNS depressants. Withdrawal symptoms have occurred following abrupt discontinuance. Keep addiction-prone individuals under careful surveillance because of their predisposition to habituation and dependence. In pregnancy, lactation or women of childbearing age, weigh potential benefit against possible hazard.

Precautions: If combined with other psychotropics or anticonvulsants, consider carefully pharmacology of agents employed. Usual precautions indicated in patients severely depressed, or with latent depression, or with suicidal tendencies. Observe usual precautions in impaired renal or hepatic function. Limit dosage to smallest effective amount

in elderly and debilitated to preclude ataxia or oversedation.

Side Effects: Drowsiness, confusion, diplopia, hypotension, changes in libido, nausea, fatigue, depression, dysarthria, jaundice, skin rash, ataxia, constipation, headache, incontinence, changes in salivation, slurred speech, tremor, vertigo, urinary retention, blurred vision. Paradoxical reactions such as acute hyperexcited states, anxiety, hallucinations, increased muscle spasticity, insomnia, rage, sleep disturbances, stimulation, have been reported; should these occur, discontinue drug. Isolated reports of neutropenia, jaundice; periodic blood counts and liver function tests advisable during long-term therapy.



Roche
LABORATORIES

Division of Hoffmann-La Roche Inc.
Nutley, New Jersey 07110

Valium® (diazepam)